CALIFORNIA ASTER Symphyotrichum chilense



Cheerful, light purple blooms, summer to fall in the San Francisco Bay Area. Butterflies and moths consume the nectar and birds eat the seed after blooms have departed. Spreads easily. Deer resistant. Also called Pacific Aster.

This plant is a member of the sunflower family.



BIODIVERSITY VALUE **MEDIUM**

BUTTERFLIES/MOTHS

BIRDS

53

Northern Mockingbirds, House Finches, California Thrashers, orioles, Cedar Waxwings, Oak Titmice, nuthatches, Chestnut-backed Chickadees, American Crows, California Scrub and Stellar's Jays, vireos, Bewick's and House Wrens, White-crowned Sparrows, Downy, Hairy and Nuttall's woodpeckers



CEDAR WAXWING



CALIFORNIA SCRUB JAY



DOWNY WOODPECKER



TYPE Perennial herb



SIZE 1.3-3.5 ft. tall / 3 ft. wide



SUN Full sun or part shade. A wandering border flower or groundcover, decorative near grassy areas. Cut back after blooms have seeded.



WATER Low



SOIL Loamy clay

Learn more at: cnps-yerbabuena.org & goldengatebirdalliance.org

Resources: Calscape, Doug Tallamy, Kathy Kramer Bringing Back the Natives
Photo credits: Noreen Weeden; Irene; Steven Smethurst; Dario Taraborelli; Andy Wilson

